arXiv:1011.6085v1 [math.GT] 28 Nov 2010

SELF-INTERSECTIONS ARE EMPIRICALLY GAUSSIAN

MOIRA CHAS

Abstract. In an orientable surface with boundary, free homotopy classes of curves on surfaces are in one to one correspondence with cyclic reduced words in a set of standard generators of the fundamental group. The combinatorial length of a class is the number of letters of the corresponding word. The self-intersection of a free homotopy class (that is, the smallest number of self-crossings of a representative of a class) can be computed in terms of the word. For each of the free homotopy classes of length twenty on the punctured torus, we compute its self-intersection number and make a histogram of how many have self-intersection 0, 1, 2..... The histogram is essentially Gaussian.

By choosing a set of free generators on the fundamental group of the punctured torus T, one can establish a one-to-one correspondence between the set of free homotopy classes of oriented curves on T and the conjugacy classes of the free group of rank two. The *combinatorial length of a conjugacy class* is the minimal number of letters (generators or inverses) required for its description.



Figure 1. The histogram showing the distribution of self-intersection numbers over all hundred and fifty million non-power classes of combinatorial length 20 in the punctured torus. The horizontal coordinate shows the self-intersection count k; the vertical coordinate shows the number of cyclic reduced words for which the self-intersection number is k.

The *self-intersection of a free homotopy class of curves* (that is, the smallest number of times in which a representative crosses itself, counted by multiplicity) can be computed using the algorithms described in [4] or [1]. We programmed these algorithms using Java [2] and performed this computation for each non-power class of combinatorial length twenty. Then we organized the output in the histogram of Figure 1 portraying proportion of classes of with each possible self-intersection number. The histogram appears clearly normal. Based on these findings there is a sequel to this note [3] that in any orientable surface with boundary and negative Euler characteristic, the distribution of self-intersection sampling classes by their combinatorial length approaches a Gaussian as the combinatorial length goes to infinity.

REFERENCES

[1] Moira Chas. Combinatorial Lie bialgebras of curves on surfaces. *Topology*, 43(3):543–568, 2004.

[2] Moira Chas. Java applet for computing the intersection and self-intersection of curves on surfaces and the goldman-turaev lie bialgebra, 2004. http://www.math.sunysb.edu/ moira/applets/.

[3] Moira Chas and Steve Lalley. Self-intersections in combinatorial topology: statistical structure. 2010.

[4] Marshall Cohen and Martin Lustig. Paths of geodesics and geometric intersection numbers. I. In *Combinatorial group theory and topology (Alta, Utah, 1984)*, volume 111 of *Ann. of Math. Stud.*, pages 479–500. Princeton Univ. Press, Princeton, NJ, 1987.